

AD-A096 641

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

A SEQUENTIAL K-GROUP RANDOM ALLOCATION METHOD WITH APPLICATIONS--ETC(U)

DEC 80 A P 50MS

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2153

NL

1 of 1  
AD 8  
(10000-21)



END  
DATE  
FILMED  
4-81  
DTIC

LEVEL II

(2)

AD A 096641

MRC Technical Summary Report #2153

A SEQUENTIAL k-GROUP RANDOM  
ALLOCATION METHOD WITH APPLICATIONS  
TO SIMULATION

Andrew P. Soms

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

DTIC  
ELECTE  
MAR 23 1981  
E

December 1980

(Received November 12, 1980)

FILE COPY

Approved for public release  
Distribution unlimited

Sponsored by

U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709

Office of Naval Research  
Arlington, Virginia 22217

81 3 19 090

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

A SEQUENTIAL k-GROUP RANDOM ALLOCATION METHOD  
WITH APPLICATIONS TO SIMULATION

Andrew P. Soms\*

Technical Summary Report #2153

December 1980

ABSTRACT

A sequential method of random allocation is given and it is shown how it can be used to estimate the observed significance levels of k-sample nonparametric tests. The sequential technique is compared to the standard random allocation technique and shown to be more efficient. An application is made to the Dunn-Bonferroni method of multiple comparisons.

AMS(MOS) Subject Classification: 62L99, 65C05

Key Words: Dunn-Bonferroni method; Nonparametric tests;  
Observed significance levels; Simulation

Work Unit No. 4 - Statistics and Probability

\*University of Wisconsin-Milwaukee, Milwaukee, Wisconsin 53201.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and the Office of Naval Research under Contract No. N00014-70-C-0321.

## SIGNIFICANCE AND EXPLANATION

A major problem in using nonparametric tests is to determine the observed level of significance. This paper gives an efficient sequential simulation method for doing this.

Simulation For	
Test	<input checked="checked" type="checkbox"/>
Method	<input type="checkbox"/>
Number of Simulations	
Initial Value	
Available for	
Dist	
A	

---

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

# A SEQUENTIAL k-GROUP RANDOM ALLOCATION METHOD WITH APPLICATIONS TO SIMULATION

Andrew P. Soms

## 1. The Sequential Allocation Method

Bebbington (1975) showed that if there were  $N$  objects (such as file cards) from which it was desired to select (without replacement here and throughout) a random sample of size  $k$  without numbering the  $N$  objects, then one could proceed sequentially by selecting the first object with probability  $k/N$  and if at the  $T^{\text{th}}$  stage  $s$  have been selected, then the  $T+1^{\text{st}}$  object is selected with probability  $(k-s)/(N-T)$ ,  $T=1,2,\dots,N-1$ .

We now state and prove the extension to an arbitrary number of groups. Suppose there are  $N$  objects and it is desired to sequentially divide them randomly into  $r$  groups of size  $k_1, k_2, \dots, k_r$ ,  $\sum_{i=1}^r k_i = N$ , i.e., each allocation has probability  $1/\binom{N}{k_1, \dots, k_r}$ . Let  $s_{1T}, \dots, s_{rT}$  be the number of objects selected for groups  $1, 2, \dots, r$  at the  $T^{\text{th}}$  stage and let  $P_{i,T+1}$  denote the selection probability for group  $i$  at the  $T+1^{\text{st}}$  stage. Then if

$$P_{i,T+1} = (k_i - s_{iT}) / (N - T), \quad T = 0, 1, \dots, N-1, \quad (1.1)$$

the selection is random. Note that  $P_{i,1} = k_i/N$  and  $\sum_{i=1}^r P_{i,T+1} = 1$ . The randomness follows immediately by noting that the probability of a particular assignment is

$$\left( \prod_{i=1}^r k_i! \right) / N! = 1 / \binom{N}{k_1, \dots, k_r}.$$

Bebbington's (1975) result is a special case of the above when  $r = 2$ .

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and the Office of Naval Research under Contract No. N00014-70-C-0321.

As an example, suppose  $r=3$ ,  $k_1=2$ ,  $k_2=2$ ,  $k_3=3$  and  $N=7$ . In order to make the sequential allocation given by (1.1) we take 7 independent random numbers  $U_i$ ,  $i = 1, 2, \dots, 7$ . Let

$$Q_{0,T} = 0 \text{ and } Q_{i,T} = \sum_{j=1}^i P_{j,T}, \quad i = 1, 2, \dots, r, \quad T = 1, 2, \dots, N.$$

Then the  $m^{\text{th}}$  object,  $m = 1, 2, \dots, N$ , is assigned to group  $n$ , where  $n$  is the unique integer such that

$$Q_{n-1,m} < U_m \leq Q_{n,m}.$$

Suppose the 7 random numbers are .79039, .01850, .99744, .81812, .93169, .22705, and .97709. The selection process is summarized in Table 1.

#### 1. Selection Process

<u>Stage</u>	<u>Random Digit</u>	<u>P<sub>1T</sub></u>	<u>P<sub>2T</sub></u>	<u>P<sub>3T</sub></u>	<u>Group Selected</u>
1	.79039	2/7	2/7	3/7	3
2	.01850	2/6	2/6	2/6	1
3	.99744	1/5	2/5	2/5	3
4	.81812	1/4	2/4	1/4	3
5	.93169	1/3	2/3	0	2
6	.22705	1/2	1/2	0	1
7	.97709	0	1	0	2

Note that if all the  $k_i$ 's are one, a random permutation is produced if we think of the group as denoting position.

## 2. Applications to Simulation

In  $k$ -sample nonparametric tests the observed significance level of the test is obtained by considering all possible partitions  $M$  of the (possibly tied) observed values or (possibly average) ranks into  $r$  groups, computing the value of the test statistic, and counting the number of times  $m$  it is equal to or greater than the observed value. The observed significance level  $\hat{\alpha}$  is then  $m/M$ . When the number of partitions is large this is prohibitive and  $\hat{\alpha}$  is estimated either by simulation (taking a large random sample of the allocations) or by asymptotics. The advantage of simulation is that one can control the accuracy of the estimate (by taking a large or small random sample) depending on the importance of the situation, unlike asymptotics which each time it is used forces one into the straight-jacket of committing a usually unknown error. Since it is (perhaps regrettably) a well known fact that different actions will be taken for close values of  $\hat{\alpha}$ , one above and the other below some fixed level (e.g., .01, .05, or .1) of the decision-maker, the use of simulation at least prevents approximating error in  $\hat{\alpha}$  to be the determining factor.

If it is decided to use simulation, then a possible procedure is to make the random assignment as described in Section 1 many times by using a computer. The commonly used method is to produce a random permutation by ordering a random sample of uniform numbers and choosing the first  $k_1$  indexes for group 1, the next  $k_2$  for group 2, and so on. If all the  $k_i$ 's are one, then this is more efficient than Section 1. However, as soon as the  $k_i$ 's

depart even moderately from 1, the method of Section 1 becomes much more efficient. As an example, if  $k_1 = k_2 = k_3 = k_4 = 10$  and it is desired to make 2000 random assignments using a UNIVAC 1110 computer, a FORTRAN program using the methods of Section 1 uses 4.71 seconds of CPU time while a FORTRAN program using the random permutation method takes 9.17 seconds.

The Appendix contains a listing of the FORTRAN subroutine RANDM that uses the theory of Section 1 to make random assignments. This may be tied in with any specific simulation problem, e.g., the case treated in Section 3.



### 3. Applications to the Dunn-Bonferroni Method of Multiple Comparisons

The D-B (Dunn-Bonferroni) method is described in Dunn (1964). Briefly, let  $Y_{ij}$ ,  $i = 1, 2, \dots, r$ ,  $j = 1, 2, \dots, n_i$ , be continuous (this assumption is not important and is removed later) random variables with distribution function  $F_i$ ,  $H_0: F_1 = F_2 = \dots = F_r$ ,  $H_a$ : for at least one pair  $(i, j)$ ,  $F_i \neq F_j$  (in the sense of producing larger or smaller values), and the test must identify which, if any, pairs are different. Denote by  $z_\alpha$  the upper  $\alpha^{\text{th}}$  point of the standard normal. The D-B test declares all those pairs  $(i, j)$ ,  $i < j$ , different for which

$$z_{ij} = |\bar{R}_i - \bar{R}_j| / \left[ \frac{(N)(N+1)}{12} \left( \frac{1}{n_i} + \frac{1}{n_j} \right) \right]^{1/2} \geq z_\alpha / (k(k-1)) , \quad (3.1)$$

where  $\bar{R}_i$  denotes the average of the ranks of the  $i^{\text{th}}$  group in the joint ranking. The nominal significance level of this procedure is  $\alpha$ . The actual significance level  $\alpha_A$  is

$$\alpha_A = P_0 \left( \max_{i < j} z_{ij} \geq z_\alpha / (k(k-1)) \right) , \quad (3.2)$$

and may be obtained by simulation based on Section 1. Table 2 gives some comparisons of nominal with actual, using Section 1 and 10,000 simulations.

## 2. Comparison of Actual to Nominal $\alpha$

<u>r</u>	<u>Common Group Size</u>	<u>Nominal <math>\alpha</math></u>	<u>Actual <math>\alpha</math></u>
3	5	.05	.037
3	10	.05	.040
3	15	.05	.043
3	30	.05	.045
3	5	.01	.0030
3	10	.01	.0077
3	15	.01	.0077
3	30	.01	.010
5	5	.05	.026
5	10	.05	.036
5	5	.01	.0030
5	10	.01	.0067

The Appendix contains a listing of the program used for Table 2. It thus appears that D-B is conservative and we can remove the conservatism by substituting for  $z_{\alpha}/(k(k-1))$   $d_{(i)}$ , where  $d_{(i)}$ ,  $i=1,2,\dots,r(r-1)/2$ , is the  $i^{\text{th}}$  largest observed values of  $z_{ij}$ ,  $i < j$ , to obtain by simulation the  $r(r-1)/2$  possible observed significance levels.

The K-S (Kruskal-Scheffé) method is also sometimes used in this situation (see, e.g., Miller, 1966, p. 166) and consists of replacing  $z_\alpha/(r(r-1))$  in (3.1) with  $h_\alpha^{1/2} = (\chi_{\alpha;r-1}^2)^{1/2}$ , where  $\chi_{\alpha;r-1}^2$  is the upper  $\alpha^{\text{th}}$  point of  $\chi^2$  with  $r-1$  degrees of freedom. The comparison of the critical constants in Table 3 shows that this is even more conservative than D-B.

### 3. Comparison of D-B and K-S Critical Constants

$r$	$z_{.05}/(r(r-1))$	$(\chi_{.05;r-1}^2)^{1/2}$	$z_{.01}/(r(r-1))$	$(\chi_{.01;r-1}^2)^{1/2}$
3	2.39	2.79	2.94	3.36
4	2.50	3.08	3.02	3.65
5	2.58	3.33	3.09	3.89
6	2.64	3.55	3.15	4.10
7	2.69	3.75	3.19	4.30
8	2.74	3.94	3.23	4.48
9	2.77	4.11	3.26	4.66
10	2.81	4.28	3.29	4.82

If the data is discrete, the D-B method can be modified as in Dunn (1964) and the random assignment done on average ranks. Thus ties present no problems in this approach.

For all practical purposes the exact D-B (use of the  $d_{(i)}$  and simulation) seems the best method to use.

# Appendix

```

C EXAMPLE OF MAIN PROGRAM FOR SEQUENTIAL RANDOM ALLOCATIONS
C NR IS THE NUMBER OF GROUPS, NSIM THE NUMBER OF SIMULATIONS, K'S THE GROUP
C SIZES, X'S THE NUMBERS TO BE ALLOCATED
DIMENSION X(1000),Z(20,100),K(20)
100 FORMAT ( )
99 READ 100, NR, NSIM
IF (NR.EQ. 0) GO TO 101
READ 100, (K(J), J=1, NR)
KK=0
DO 1 J=1, NR
1 KK=KK+K(J)
DO 203 I=1, KK
203 X(I)=1
DO 2 II=1, NSIM
CALL RANDM(KK, K, KK, X, Z)
2 CONTINUE
GO TO 99
101 STOP
END
#FOR, IS, SUB1
SUBROUTINE RANDM(NR, K, KK, X, Z)
C NR NUMBER OF GROUPS, K ARRAY OF GROUP SIZES, KK NUMBER OF ELEMENTS, X ARRAY
C OF KK ELEMENTS, Z RANDOM ALLOCATION OF X
DIMENSION X(1000), Z(20, 100), NSC(20), GC(20), QC(20), U(1000), K(20)
DO 3 I2=1, NR
3 NSC(I2)=0
DO 333 I2=1, KK
333 U(I2)=RANUN(X)
DO 4 I3=1, NR
4 GC(I3)=0
DO 5 II=1, MAXI
DO 5 III=1, II
5 GC(II)=GC(II)+U(III)
DO 6 I4=1, KK
U(I4)=(KK-I4+1)*U(I4)
IF (U(I4).LE.GC(1)) GO TO 61
DO 7 I5=2, MAXI
7 IF (U(I4).LE.GC(I5)) GO TO 62
IF (U(I4).GT.GC(NR-1)) INDEX=NR
GO TO 64
61 INDEX=1
GO TO 64
62 INDEX=I5
64 NSC(INDEX)=NSC(INDEX)+1
NN=NSC(INDEX)
Z(INDEX, NN)=X(I4)
C UPDATE
DO 8 I6=1, NR
8 GC(I6)=0
DO 9 II=1, MAXI
DO 9 III=1, II
9 GC(II)=GC(II)+U(III)
6 CONTINUE
RETURN
END

```

C D-R BY SIMULATION  
C NR IS THE NUMBER OF GROUPS, NSIM THE NUMBER OF SIMULATIONS, ZAL IS THE POINT  
C ~~FOR WHICH THE PROBABILITY OF THE MAXIMUM OF ALL THE ABSOLUTE VALUES OF~~  
C STANDARDIZED RANK AVERAGE DIFFERENCES EQUALLING OR EXCEEDING IT IS TO BE  
C CALCULATED, K'S ARE THE GROUP SIZES

```

DIMENSION IFLAG(20,20),R(20)
DIMENSION X(1000),Z(20,100),K(20)
100 FORMAT ( )
99 READ 100, NR, NSIM
  IF (NR.EQ. 0) GO TO 101
  MAXI=NR-1
READ 100, ZAL
  READ 100, (K(J), J=1, NR)
  KK=0
  DO 1 J=1, NR
    1 KK=KK+K(J)
  DO 203 I=1, KK
203 X(I)=1
    CON=KK*(KK+1)*ZAL**2
    COUNT=0
    DO 2 I1=1, NSIM
      CALL RANDM(NR, K, KK, X, Z)
      DO 20 J1=1, NR
        R(J1)=0
        NUJP=K(J1)
        DO 20 J2=1, NUJP
20 R(J1)=R(I1)+Z(J1, J2)
          DO 21 J1=1, MAXI
            LLIM=J1+1
            DO 21 J2=LLIM, NR
              IFLAG(J1, J2)=0
21 IF (12*(K(J2)*R(J1)-K(J1)*R(J2))**2.GE.K(J1)*K(J2)*CON*(K(J1)+K(J2
1))) IFLAG(J1, J2)=1
              IPRD=0
              DO 22 J1=1, MAXI
                LLIM=J1+1
                DO 22 J2=LLIM, NR
22 IPRD=IPRD+IFLAG(J1, J2)
                IF (IPRD.GE.1) COUNT=COUNT+1
2 CONTINUE
      PR=COUNT/NSIM
      PRINT 100, (K(I), I=1, NR)
      PRINT 100, ZAL, NSIM, PR
      GO TO 99
101 STOP
  END

```

### References

- Bebbington, A. C. (1975), "A Simple Method of Drawing a Sample Without Replacement", Applied Statistics, 24, 136.
- Dunn, Olive J. (1964), "Multiple Comparisons Using Rank Sums", Technometrics, 6, 241-52.
- Miller, Rupert J. (1966), Simultaneous Statistical Inference, New York: McGraw-Hill Book Company.

14) MRZ-TSR-1413

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TSR# 2153	2. GOVT ACCESSION NO. AD-A096642	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Sequential k-Group Random Allocation Method with Applications to Simulation,		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) 10) Andrew P. Soms		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) 15) DAAG29-80-C-0041, N00014-70-C-0321
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit No. 4 (Statistics & Probability)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9) Technical Summary Repts		12. REPORT DATE 11) December 1980
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U.S. Army Research Office and Office of Naval Research P.O. Box 12211 Arlington, Virginia 22217 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Dunn-Bonferroni method, Nonparametric tests, Observed significance levels, Simulation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A sequential method of random allocation is given and it is shown how it can be used to estimate the observed significance levels of k-sample nonparametric tests. The sequential technique is compared to the standard random allocation technique and shown to be more efficient. An application is made to the Dunn-Bonferroni method of multiple comparisons.		

FILM  
4-1